

**CARN Examination Blueprint  
Ideal Distribution of 150-Item Test**

→ Domains of Practice  Patient problems ↓	<b>1</b> Perform a biopsychosocial SUD assessment	<b>2</b> Assess acute care needs of SUD patients	<b>3</b> Develop and implement an individualized plan of care	<b>4</b> Educate and promote behavioral change	<b>5</b> Care mgmt., treatment, and evaluation, throughout the recovery continuum (including special populations)	<b>TOTAL</b>
<b>A</b> Opioid use disorder	5-6	5-6	7-8	5-6	13-14	<b>25%</b> <b>(37-39)</b>
<b>B</b> Alcohol use disorders	5-6	5-6	7-8	5-6	13-14	<b>25%</b> <b>(37-39)</b>
<b>C</b> Medication misuse (sedative/hypnotics/anxiolytics, gabapentinoids, etc)	3-4	3-4	4-5	3-4	8-9	<b>15%</b> <b>(22-24)</b>
<b>D</b> Co-occurring psychiatric/comorbid medical disorders (process addictions, infectious diseases, and mental health disorders)	3-4	3-4	4-5	3-4	8-9	<b>15%</b> <b>(22-24)</b>
<b>E</b> Stimulant use disorders (cocaine, amphetamines, caffeine)	2-3	2-3	3-4	2-3	5-6	<b>10%</b> <b>(14-16)</b>
<b>F</b> Cannabinoids and other hallucinogens	1-2	1-2	1-2	1-2	2-3	<b>5%</b> <b>(7-9)</b>
<b>G</b> Tobacco use disorder (e.g., vaping, nicotine)	1-2	1-2	1-2	1-2	2-3	<b>5%</b> <b>(7-9)</b>
<b>Total</b>	<b>15%</b> <b>(22-24)</b>	<b>15%</b> <b>(22-24)</b>	<b>20%</b> <b>(29-31)</b>	<b>15%</b> <b>(22-24)</b>	<b>35%</b> <b>(52-54)</b>	<b>100%</b> <b>(150)</b>