

CARN Test Specifications (Blueprint)
Ideal Distribution of 150 Items

→ Domain of Practice Patient problem ↓	1 Perform a biopsychosocial SUD assessment	2 Assess acute care needs of SUD patients	3 Develop and implement an individualized plan of care	4 Educate and promote behavioral change	5 Care mgmt., treatment, and evaluation, throughout the recovery continuum (including special populations)	TOTAL
A Opioid use disorder	5-6	5-6	7-8	5-6	13-14	27% (40-42)
B Alcohol use disorders	5-6	5-6	7-8	5-6	13-14	25% (37-39)
C Medication misuse (sedative/hypnotics/anxiolytics, gabapentinoids, etc)	3-4	3-4	4-5	3-4	8-9	10% (15-17)
D Co-occurring psychiatric/comorbid medical disorders (process addictions, infectious diseases, and mental health disorders)	3-4	3-4	4-5	3-4	8-9	12% (19-21)
E Stimulant use disorders (cocaine, amphetamines, caffeine)	2-3	2-3	3-4	2-3	5-6	15% (22-24)
F Cannabinoids and other hallucinogens	1-2	1-2	1-2	1-2	2-3	6% (9-11)
G Tobacco use disorder (e.g., vaping, nicotine)	1-2	1-2	1-2	1-2	2-3	5% (7-9)
Total	15% (22-24)	20% (29-31)	15% (22-24)	15% (22-24)	35% (52-54)	100% (150)